The Growing Stronger Program

The Growing Stronger Tool Kit

A Program Leader's Guide to Conducting

Strength Training Programs for Older Adults

Rebecca Seguin, MS, PhD, CSCS

Cornell University
Ithaca, NY

Miriam E. Nelson, PhD

University of New Hampshire
Durham, NH

Jill FitzGerald, PT, DPT, GCS, CSCS, CEEAA

Saint Louis University
Saint Louis, MO

Kelly Hawthorne, PT, DPT, GCS, CEEAA

Saint Louis University
Saint Louis, MO

Caution

The Growing Stronger Program is an exercise program based upon extensive scientific research. The Growing Stronger Tool Kit contains instructions and safety precautions for community exercise leaders. It is essential that you read the entire Tool Kit carefully before implementing the program. Some exercises outlined in the program are not appropriate for individuals with unstable medical conditions or serious musculoskeletal problems.

This Tool Kit is not intended to replace the services of a healthcare provider who knows the participants of your program personally. As a fitness professional, never give medical advice to a participant in your program. Always refer the individual to his/her healthcare provider. Individuals who participate in the Growing Stronger Program are strongly encouraged to talk with their doctors before starting this program.

Every effort has been made to ensure that the information contained in this Tool Kit is complete and accurate. However, neither the authors nor Cornell University is engaged in rendering medical advice or services to individual participants in the Growing Stronger Program. The ideas, procedures, and suggestions contained in this Tool Kit are not intended as a substitute for consulting with a physician. All matters regarding a participant's health require medical supervision. Neither the authors nor Cornell University shall be liable or responsible for any loss, injury, or damage allegedly arising from any information or suggestion in this Tool Kit. The opinions expressed in this Tool Kit represent the personal views of the authors and not those of Cornell University.

Mission

To increase the number of older adults participating in safe and effective strength training programs throughout the United States and abroad.

Objectives

™ To provide the necessary educational materials and training to Program Leaders to help them successfully implement and maintain the Growing Stronger Program.

™ To facilitate the implementation of the Growing Stronger Program in community centers, local health and fitness facilities, hospitals and clinics, cooperative extension services, businesses, and any other place where older adults come together.

™ To improve the health and well-being of older adults throughout the United States and abroad by increasing access to structured, safe, and effective strength training programs.

showed that strength training increases bone density and reduces the risk for fractures among women aged 50 to 70.[1]

Proper Weight Maintenance

Strength training is crucial to weight control, because individuals who have more muscle mass have a higher metabolic rate. Muscle is active tissue that consumes calories, whereas stored fat uses very little energy. Strength training can provide up to a 15% increase in metabolic rate, which is enormously helpful for weight loss and long-term weight control.[13]

Improved Glucose Control

More than 16 million Americans have type 2 diabetes, a disease in which the body's cells do not properly respond to insulin, thereby causing blood glucose levels to rise.[14] Strength training can help prevent and control diabetes by increasing the body's sensitivity to insulin, thereby improving glucose control.[15]

© Cornell University

© Cornell University

©

© Cornell University

™ Allow about 3-4 weeks between the informational meeting and the start date of the program to obtain participant medical clearance

Part F Scheduling the program

Each program site can run the Growing Stronger Program according to the schedule that works best for the community. Generally, the program is offered two days per week, but it can be run three days per week if desired. The only scheduling requirement is that the exercise sessions never meet on consecutive days of the week because participants will need at least one day off between sessions. Session length will be approximately one hour; this should be kept in mind when schedules are being made. We have found that 12-, 14-, and 16-week sessions with scheduled breaks work well for programs in community settings. For instance, you may want to run a 12-week program followed by a 4-week break, and then begin another 12-week program. Another option is a 14-week session with a 2-week break. As the program gets under way, you and the manager of the program site should discuss whether the program schedule is working well and make adjustments at the end of the first session.

© Cornell University

Part B Participant's medical history

Medical History and Current Health Survey

Name_____

Please read the following list carefully and circle Yes or No as it applies to your medical history and current health. Please include any additional information and conditions for which you are receiving medical care.

Medical History		
Aneurysm	Yes	No
Arthritis (Rheumatoid or Osteoarthritis)	Yes	No
Asthma	Yes	No
Back Pain	Yes	No
High Blood Pressure (Last reading /)	Yes	No
Low Blood Pressure (Last reading /)	Yes	No
Bone Fractures	Yes	No
Cancer (Please provide type and treatment)	Yes	No
High Cholesterol (Last reading /)	Yes	No

TTdY

© Cornell University

The

Part C PAR-Q (Physical Activity Readiness Questionnaire)

Source: Physical Activity Readiness Questionnaire (PAR-Q), c. 2002. Reprinted with permission from the Canadian Society for Exercise Physiology. http://www.csep.ca/forms.asp

© Cornell University

Part D Participant and physician consent

Participant Consent

I have voluntarily enrolled in a program of progressive exercise. The program is designed to place a gradually increased workload on the heart, lungs, muscles, and bones to help improve their function. I understand that participation in such a program may be associated with some risks. These risks may include but are not limited to muscle soreness, fainting, disorders of heart beat, abnormal blood pressure, and in very rare instances, heart attack. To the best of my knowledge I do not have any limiting physical conditions or disability that would preclude an exercise program. Effort will be made to minimize any risks to me by a pre-exercise assessment and a medical screening. I release everyone who has designed, promoted, or conducted the Growing Stronger Program from all claims or liabilities whatsoever resulting from my participation in this program. I assume all risks and responsibility for any injury, damage, or any other adverse event that may result from my participation in this program.

Before I begin this program I understand that a pre-exercise assessment and physician screening consent form may be required. I

© Cornell University

"Dear Doctor" Letter

Date: _____

Dear Dr._____,

Your patient_____, is interested in participating in the Growing Stronger Program at _____. This moderate intensity, progressive exercise program includes strength and balance training and is designed to improve muscle strength, dynamic balance, and flexibility.

This program is based upon the results of strength training studies in older adults conducted by scientists at the John Hancock Research Center on Physical Activity, Nutrition, and Obesity Prevention at the Gerald J. and Dorothy R. Friedman School of Nutrition Science and Policy at Tufts University, Boston, MA. Scientists and exercise physiologists at Tufts University have designed this exercise program especially for older adults, and Program Leaders in your community are implementing the program. Your patient will be required to complete a Medical History Questionnaire and provide Informed Consent prior to participation in this exercise program.

Please complete and sign the enclosed Physician Authorization Form. If you have any questions or would like to discuss your patient's participation in this program in further detail, please call _____(name) at:_____(phone number).

Sincerely,

_____

© Cornell University

Physician Authorization Form

Patient Name:_____

Address:_____

_____

Phone Number:_____Date of Birth:_____

Date of Last Exam:_____

Height:_____Weight:_____Pulse:_____BP:_____ Other:____

_____

Medical Conditions:_____

Medications:_____

_____

Special Considerations: _____

_____

_____

_____Yes, my patient can participate.

_____ No, my patient cannot participate at this time due to his/her medical conditions and health status.

Physician's Signature:

_____

Print Name:

_____

Address:

_____

Phone Number: _____Fax Number: _____

© Cornell University

Chapter 4: The

© Cornell University

Balance is the ability to maintain equilibrium when disrupted by an internal or external force. Many physical factors that affect balance include vision, hearing, muscle strength, flexibility, and posture. Certain medical

© Cornell University

© Cornell

point at which you should encourage a small increase in the amount of weight being lifted or encourage participant to engage in higher level activity (as guided by the video).

Intensity

It is imperativof

Breathing

Participants should breathe through the mouth continuously and regularly throughout the exercises. This can be done in one of two ways. First, participants may count out loud to keep the pace of the exercises. Talking (counting) ensures that participants are not holding their breath. The second method entails inhaling before the lift, exhaling through the mouth while lifting, often referred to as "exhale during the exertion," and inhaling through the nose during the lowering phase. This method is more difficult to master initially but is very effective in reducing the monotony of counting in advanced or long-term groups.

It is important for participants to start out at an easy or moderate level for all of these exercises. When the weight is light, the participant can safely learn the unique moves of each exercise and learn how to breathe properly. After mastering proper exercise technique, then the participants can start to progress and meet the appropriate intensity for an effective workout.

Part B Program safety

With any type of exercise, there is some degree of risk; however, the potential benefits are usually far greater than the chance of adverse events. In fact, the risks are generally lower with strength training than for many other sports and activities.[16] It is extremely important that close attention be paid to the safety of participants at all times. This includes exercise technique and progression safety as well as general equipment safety. You should always be conscious of the technique and progression of each participant—correcting, demonstrating, and guiding proper form whenever

© Cornell University

necessary. Discourage sudden or quick movements, particularly when using weight equipment. Movements to avoid include swinging of the legs, swinging out of the arms, and walking around the exercise area while wearing ankle weights.

While encouraging progression is vital, you must always discourage participants from lifting any weight that is too heavy or from doing any exercise that causes pain or discomfort. Take a conservative approach until you have determined the participant's level of ability.

Drinking water should be readily available in the exercise area. Encourage the participants to drink water by taking drinking breaks throughout the exercise session.

Encourage participants to breathe normally through their mouths.ir

As a Program Leader for older adults, it is your job to be the eyes and ears of the physician. If there are any changes in the exercise participant's condition, recommend that he/she consult his/her health care provider.

Unstable medical conditions or severe trauma in the past six months are contraindications for exercise. Be sure that all potential exercisers are under a doctor's care. Every week before beginning to exercise, ask each exercise participant if he/she has had any changes in his/her health or medication.

Medical issues should always be referred to the participant's physician or health care provider. **Program Leaders should never give medical advice to participants.**

Pay particular attention to the back, shoulders, and knees. Emphasize the importance of protecting the lower back by bending down from the knees and not bending at the waist. Participants should be reminded to continually check the position of their knee joints when standing. The knee joints should always be slightly bent or "loose" and "soft," which means not pushed back, locked, or excessively straight. The elbows should always be bent when lifting the arms overhead. If a participant reports arthritis, bursitis, or rotator cuff pain, replace overhead movements with lower movements in the pain-free range.

To ensure general safety throughout the exercise sessions, you should make sure that ankle weights and dumbbells are always properly stored when not in use. It is also important that the exercise room and any adjoining area be well lit and free of clutter or spills. Any defective or damaged equipment – including chairs – should be removed immediately

© Cornell University

until repaired. Coats, sweaters, purses, and personal items must be hung up or put in a designated area. They may not be kept underneath or near the participant's chair.

Participants should be encouraged to wear appropriate clothing and shoes. Comfortable clothes that are easy to move around in are recommended. Skirts may be worn; however, pants or shorts are encouraged. Proper footwear is reqQ/>p-82 0 uTj -0.008 Tc44(u)8.3re cBox4 uTj -0.008 T

Chapter 5: Keeping track & participant assessments

Part A

© Cornell University

[redacted]t attendance sheet

[redacted], you are encouraged to keep a record of participant [redacted]cise session. This will help you monitor the success

[redacted]rticipant Attendance Sheet

[redacted]_____**Site**_____

[redacted]_____

[redacted]	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/

am © Cornell University

The

Date			
Warm Up			

© Cornell University

Part D Participant evaluation

At the completion of a 12-week session of a Growing Stronger Program, participants are strongly encouraged to fill out the following evaluation form.

Participant Evaluation

Participe b 0.004 Tw 12 -0 0 12

Chair Stand Test

Purpose:

To measure lower-body strength.

Equipment:

Stopwatch and straight-back or folding chair with a seat height of 17 in (43.18 cm). Chair is placed against a wall to prevent slipping.

©Cornell University

The

Arm Curl Test

Purpose:

To measure upper-8.3(e)0.6(a)0.6004 Tc 004 Te05 0 Td (-)2.3Tj /P <</MCI4 Tw 14.04 -0 0

The

©Cornell Universit

2-Minute Step Test

Purpose:

To assess aerobic endurance (th 2 201.5t is 3.6(() .2(o)8ltr(ur)3.6(a)1.2(o)8tabi6.709 0 Td ( )T

knee reaches the target. When the proper knee height cannot be maintained, ask the participant to slow down, or to stop until he/she can regain the proper form, but keep the time running.

Scoring:

The score is the number of full steps in 2 minutes; that is, the number of times the right knee reaches the proper height. Administer only one trial on test day. However, for maximal scoring accuracy, have participant practice the test (stepping in place for 2 minutes) on a day prior to the test.

HowevengP

©Cornell University

2-Minute Step Test

Adapted from Rikli & Jones 1999. Reprinted, by permission, from R.E. Rikli & C.J. Jones, 2001, Senior Fitness Test Manual, (Champaign, IL: Human Kinetics), pages 61-74 and 125-131.

Chair Sit-and-Reach Test

Purpose:

To assess lowerRandt

The Growing Stronger

Chair Sit-and-Reach Test

Adapted from Rikli & Jones 1999. Reprinted, by permission, from R.E. Rikli & C.J. Jones, 2001, Senior Fitness Test Manual, (Champaign, IL: Human Kinetics), pages 61-74 and 125-131.

©Cornell University

if the middle fingers overlap. Always measure the distance from the tip of one middle finger to the tip of the other, regardless of their alignment behind the back.

Safetyprecautions:

Safetyprecautions:

When administering the 8-foot up-and-go test, stand between the chair and cone in order to assist the participant in case he/she loses his/her bala

6-Minute Walk Test (to be completed if 2-min step test is not used)

Purpose: To assess aerobic endurance.

Equipment:

Long measuring tape, two stopwatches, four cones (or similar markers), masking tape, magic marker, 12 to 15 Popsicle sticks per person (or index cards and pencils to keep track of laps walked), chairs for waiting partners and for walkers who need to rest, and name tags.

Setup:

Mark off in 5-yard segments a flat, 50-yard rectangle (20 yards by 5 yards). The inside corners of the measured distance should be marked with cones, and the 5-yard lines marked with masking tape or chalk. (In metric units, this is a 45.7-meter course marked off in 4.57-meter segments).

Procedure:

If the 6-

©Cornell University

track of the distance walked, partners give Popsicle sticks (or similar objects) to participants each time they complete a lap. Or partners can mark a scorecard each time a lap is completed, using the "picket fence" system.

The timer should move to the inside of the marked area after everyone has started. To assist with pacing, the remaining time should be called out when walkers are about half done, and when about 2 minutes are left. Participants can stop and rest on the chairs provided, but the time keeps running. The tester should encourage participants a few times by saying, "you're doing well" and "keep up the good work." When a participant's 6 minutes has elapsed, the tester asks him/her to stop, move to the right (across from the nearest 5-yard marker), and slowly step in place for a minute to cool down.

Scoring:

Record the scores when all the walkers have been stopped. Each Popsicle stick (or mark on a card) represents 50 yards. For example, if a person has04 Twm siclaicipa), and J 0.0

©Cornell

The

Arm Curl Test (Women)

Percentile rank	60-64	65-69	70-74	75-79	80-84	85-89	90-94
95	24	22	22	21	20	18	17
90	22	21	20	20	18	17	16
85	21	20	19	19	17		
80	20	19	18	18			
75	19	18	17	17			

©Cornell University

Chair Sit-and-Reach Test (Women)

Percentile rank	60-64	65-69	70-74	75-79	80-84	85-89	90-94
95	8.7	7.9	7.5	7.4	6.6	6.0	4.9
90	7.2	6.6	6.1	6.1	5.2	4.6	3.4
85	6.3	5.7	5.2	5.2	4.3	3.7	2.5
80	5.5	5.0	4.5	4.4	3.6	3.0	1.7
75	4.8	4.4	3.9	3.7	3.0	2.4	1.0
70	4.2	3.9	3.3	3.2	2.4	1.8	0.4
65	3.7	3.4	2.8	2.7	1.9	1.3	-0.1
60	3.1	2.9	2.3	2.1	1.4	0.8	-0.7
55	2.6	2.5	1.9	1.7	1.0	0.4	-1.2
50	2.1	2.0	1.4	1.2	0.5	-0.1	-1.7
45	1.6	1.5	0.9	0.7	0.0	-0.6	-2.2
40	1.1	1.1	0.5	0.2	-0.4	-1.0	-2.7
35	0.5	0.6	0.0	-0.3	-0.9	-1.5	-3.3
30	0.0	0.1	-0.5	-0.8	-1.4	-2.0	-3.8
25	-0.6	-0.4	-1.1	-1.3	-2.0	-2.6	-4.4
20	-1.3	-1.0	-1.7	-2.0	-2.6	-3.2	-5.1
15	-2.1	-1.7	-2.4	-2.8	-3.3	-3.9	-5.9
10							

©Cornell University

8-Foot Up-and-Go Test (Women)

Percentile rank	60-64	65-69	70-74	75-79	80-84	85-89	90-94
95	3.2	3.6	3.8	4.0	4.0	4.5	5.0
90	3.7	4.1	4.0	4.3	4.4	4.7	5.3
85	4.0	4.4	4.3	4.6	4.9	5.3	6.1
80	4.2	4.6	4.7	5.0	5.4	5.8	6.7
75	4.4	4.8	4.9	5.2	5.7	6.2	7.3
70	4.6	5.0	5.2	5.5	6.1	6.6	7.7
65	4.7	5.1	5.4	5.7	6.3	6.9	8.2
60	4.9	5.3	5.6	5.9	6.7	7.3	8.6
55							

©Cornell University

6-Minute Walk Test (Women)

Percentile rank	60-64	65-69	70-74	75-79	80-84	85-89	90-94

©Cornell University

©Cornell Univer

©Cornell University

the exercises (but don't force it), smile when it's over – GENUINE smiles make people happy, and happy people means healthy people!

Learn names…and more

Knowing a person's name is a good preliminary way to make him/her feel important and to show that you really care. Fine, you say, but what if I'm just really bad with names? Here are some suggestions for learning names:

- ™ Ask the participants for a current photo when they register, and practice learning their names at home.
- ™ Have them wear nametags the first class, but be careful not to become dependent on the nametag.
- ™ Do an "ice-breaker" the first class, and have everyone tell [illegible]

Part E Communication

An important group dynamic to foster is that of open, productive communication.

When the class is4d3.5(t r)Tj E0ion 9n14.5e g(on)8. /TT1 1 Tf 9.74.5(t)-4 sc n /5(s)5.4(s)-3

"difficult" participant; someone who talks constantly, interrupts as you teach the class, protests the activities, or someone with whom you just have trouble getting along.

The worst-case scenario would be for one person to spoil the class for everyone involved, and as the instructor, you should try to prevent this from happening. Try to use the power you have as a strong leader to hear everyone's voice and encourage everyone to be an active participant in the program. Here are a few suggestions for handling a problem presented by one participant that could damage the effectiveness of the program.

- The first step is simply in remembering to **stay on task**. Try not to get caught up in difficult situations; just move on if possible.
  - If necessary, remind others to stay on task. No one wants to waste time. Hopefully this reminder with be enough to help the group move on.
  - Practice **REDIRECTING**

comments?" OR "Remember everyone, right now we're doing stretches."

™If #1 doesn't work, try **changing the subject** in a gentle way.

™ If someone is talking about something that is simply inappropriate for the class setting and she doesn't respond to #1 or 2, **tell him/her the comment or subject matter is inappropriate for the class**.

™Have a **private meeting** with the individual.

Leadership Compass

The Leadership Compass helps us to assess and understand our personal style of leadership. Most times, the Compass is used by groups who need to work on projects together to accomplish specific goals. The Compass helps them become aware of their similarities and differences, as well as what they need to change to be able to work more effectively with each other. In the case of the Growing Stronger Program, we believe that the Compass can help an individual become a better leader by allowing him/her to look at his/her natural preferences when leading a group, giving orders or suggestions, or listening to feedback and advice. Realizing how you interact with others according to your style, or "direction" on the compass, may enable you to think of ways to be a more effective leader, by capitalizing



©Cornell University

Potential Pitfalls

You may face these “pitfalls,” or personal challenges when teaching the Growing Stronger Program. Pick out those you believe you may face while

ł3.7(aation >>(3.7(aatio>-0 0 14.04.5(r)12./Artifact <<2 1.22n4 Tc 0.004 Tw 273.385 Tw 0.248

Using the Leadership Compass to Understand A Simple Task

1. Setting up the room before every class (taking out chairs and weights for the

The

©Cornell Universit

Chapter 7: Physical activity

The benefits of living a physically active lifestyle are amazing. As a Program Leader, it is important for you to understand what physical activity is and to be able

r

he/she should start slowly, with only five to ten minutes of activity, and increase the length of time as he/she adapts to the new activity.

Intensity

Intensity refers to how hard the body is working to perform a given physical activity. Intensity level is determined based on the type and the duration of an activity. Activities can be classified into one of three intensity levels: leisurely, moderate, or vigorous, typically classified by how much energy it takes to complete the activity. Leisurely activities can be described as "unhurried" and feel "easy" for a participant to complete. Leisurely activities typically last for less than 10 minutes. Strolling to the mailbox, golfing with a cart, and light gardening are examples of leisurely activities. Moderate activities consist of movements that use large muscle groups and are usually performed for ten minutes or more. Walking 1 mile in 15 minutes, bicycling 5 miles in 30 minutes, and heavy gardening are examples of moderate activities. Vigorous activities are rhythmic repetitive activities that use larger muscle groups. These activities are more intense and often last for 30 minutes or longer, though one can also complete a short burst of vigorous activity. Examples of vigorous activities are running 3 miles in 30 minutes or swimming one-half mile or more, uninterrupted. The following are some additional examples of leisurely, moderate, and vigorous physical activities:

- **Leisurely activities:** light cleaning, light gardening/yard work, hand washing a car, raking leaves, walking to a destination (less than 10 minutes), taking the stairs, golfing with a cart

- **Moderate activities:** heavy gardening, bicycling 5 miles in 30 minutes, raking leaves for 30 minutes, walking 2 miles in 30 minutes, doing water aerobics for 30 minutes
- **Vigorous activities:** swimming laps for 20 minutes, bicycling 4 miles in 15 minutes, jumping rope for 15 minutes, running 3 miles in 30 minutes, shoveling snow for 15 minutes, stair climbing for 15 minutes

The current level of fitness of each participant will play a role in determining the intensity level of a particular activity as well. A one-mile walk in 20 minutes may be moderate for a participant who has been completely sedentary, but leisurely for one who has been jogging on a regular basis. Therefore, it is important to teach participants to "listen to their body" in order to exercise at an intensity that is appropriate for them. There are three different methods you can use to help your participants determine the intensity level of their activities. They are the "talk test", the rate of perceived exertion (RPE) scale, and measured heart rate.

Talk Test

The "talk test" is a simple way to gauge the intensity level of a given activity. The test estimates a person's intensity level by determining how well he/she is able to talk while engaged in an activity. If a participant can sing while performing an activity, the activity would be considered leisurely. If he/she can carry on a conversation reasonably well (but not sing), it would be considered moderate. If a participant is too winded to have a conversation during activity, then the activity is vigorous.

Rate of Perceived Exertion (RPE)

The RPE scale is another method of gauging the intensity of physical activity.

RPE is based on one's perception of effort and is generally an accurate determinate

© Cornell University

The

i A Word of Caution *i*

Other factors affect heart rate besides activity. If any of the following situations applies to one of your participants, he/she should NOT use heart rate as a measure of intensity. Instead, he/she should be instructed

Type

Type refers to the kind of activity that will be performed (biking, gardening, etc.). A single activity is not best for everyone. Each participant should choose an activity based on what he/she enjoys, what he/she has access to, and what his/her body can handle. Encourage your participants to try a range of different activities.

Staying Active for Life

Once you have given your participants the tools they need to understand why and how they should become physically active, it is important to help them learn how to sustain an active lifestyle. Below are some tips they can use to help them continue the good habits they have learned throughout your program.

S.M.A.R.T. Goals When taking on a challenge, such as staying physically active, it is a good idea to set goals. Having realistic, defined goals helps to keep one focused on and motivated for the challenge that lies ahead. Before starting a new physical activity program, encourage your participants to set both short and long- term goals related to physical activity. These goals should be S-M-A-R-T:

S – specific…M – measurable…A – attainable…R – relevant…T – time-based

For instance: A **specific** short-term goal may be to start a program of physical activity three days per week; the long-term goal may be easing the symptoms of arthritis, improving balance, or controlling weight. This goal is easily **measurable**: Have you or have you not begun the program? Indeed, this is an **attainable** goal, as long as the doctor approves. And this goal is certainly **relevant** to living a long, healthy life. Keep the goal on a schedule: I will start within the next 2-3 days and continue for one month.

By setting S.M.A.R.T. goals, your participants will always be in the process of succeeding. These mini-successes will keep them excited and motivated to continue with their program of physical activity.

Identify Local Public Facilities

Many fitness and community centers have specific programs geared toward older adults. These programs are often not well advertised, making it difficult for your participants to learn about them. Inquire about these programs for your participants, and keep them informed as you discover new ones.

Encourage your participants to inquire on their own as well. Keeping your participants aware of these programs will help to increase their physical activity levels outside of their participation in their strength training classes. Local public facility options include (but are not limited to):

- Gyms/fitness centers
- YMCA/YWCA
- Senior/community centers
- Church groups
-

The

A: No, we suggest that participants take the day off from exercise if they have had flu symptoms or feel sick. Their bodies need time to rest and to recover. A few days off from exercise will not diminish any of the benefits they have gained from exercising.

Q: A participant has missed several exercise sessions. Should he/she use the same amount of weight that he/she was using before?
A: Have him/her start back slowly by decreasing the amount of resistance -- approximately half the weight he/she was lifting before she stopped. He/she will quickly return in a couple of weeks to where he/she left off.

Q: A participant has pain in her right shoulder. Can she just exercise the left arm?
A: First, make sure the participant determines the source of the pain and possible treatments by talking with her health care provider. If she has already done this and has arthritis or bursitis or has had a radical mastectomy or a stroke, have the participant exercise only in a range of motion that does not cause pain, or try substituting a different exercise that uses similar muscles. We are all slightly stronger on one side of our body, called the dominant side. Strength imbalances – when one side is much weaker than the other – can lead to injury. Have the participant try to keep the weights equal or within 1-2 lbs (1 kg) of each other.

© Cornell University

ACKNOWLEDGMENTS

First and foremost, we wish to extend our deepest gratitude to all of the participants in our research studies, our community strength training programs, and to the amazing leaders who teach those community program classes to thousands of women throughout the U.S. and beyond.

We wish to thank our colleagues for their continued support as well as their significant contributions to our common goal of improving people's lives through sound nutrition and physical activity. In particular, many thanks to David Buchner, Jacqueline Epping, M.Ed., and Rina Bloch, MD, who were leaders on the original leadership team that developed and tested the Growing Stronger Program.

As well, we are deeply grateful to Meagan McKinnon for her hard work on the physical activity chapter of this Tool Kit and other aspects of this project, and to Melinda Coolidge for her contribution to the chapter on leadership.

In addition, we would like to thank the students of Geriatric Seminar class of 2017 at Saint Louis University Program in Physical Therapy for the endless hours they dedicated to the enhancement of this program and production of the videos.

With gratitude and strength,

Rebecca A Seguin, PhD, CSCS
Jill M FitzGerald, PT, DPT, GCS, CSCS, CEEAA
Kelly Hawthorne, PT, DPT, GCA, CEEAA

References

1. Nelson M, Fiatarone M, Morganti C, Trice I, Greenberg R, Evans W. Effects of high-intensity strength training on multiple risk-factors for osteoporotic fractures - a randomized controlled trial. JAMA-Journal of the American Medical Association. 1994;272(24):1909-1914.
2. Castaneda C, Layne J, Munoz-Orians L, et al. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. Diabetes Care. 2002;25(12):2335-2341.
3. Beniamini Y, Rubenstein JJ, Faigenbaum AD, Lichtenstein AH, Crim MC. High-intensity strength training of patients enrolled in an outpatient cardiac rehabilitation program. Journal of Cardiopulmonary Rehabilitation and Prevention. 1999;19(1):8-17.
4. Layne J, Nelson M. The effects of progressive resistance training on bone density: a review. Medicine and Science in Sports and Exercise. 1999;31(1):25-30.
5. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. Journal of Rheumatology. 2001;28(7):1655-1665.
6. Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. Journals of Gerontology Series A-Biological Sciences and Medical Sciences. 1997;52(1):M27-M35.
7. Singh N, Clements K, Fiatarone M. Sleep, sleep deprivation, and daytime activities - a randomized controlled trial of the effect of exercise on sleep. Sleep. 1997;20(2):95-101.
8. Roubenoff R, Hughes V. Sarcopenia: current concepts. Journals of Gerontology Series A-Biological Sciences and Medical Sciences. 2000;55(12):M716-724.
9. -9.

© Cornell University

15.

© Cornell University